

NEUROBEHAVIOURAL SCREENING OF *MELALEUCA LEUCADENDRA* IN ANIMAL MODELS

Ashutosh Mishra*, Manvendra Shukla

Daksh Institute of Pharmaceutical Science, Chattarpur, M.P, India.

ABSTRACT

Melaleuca Leucadendra is common plant in India is considered carminative, stimulant and emmenagogue. In the present study, ethanol extract of bark of *Melaleuca Leucadendra* have been evaluated for Neurobehavioural study The CNS activity was assayed in several experimental models for Epilepsy i.e. Threshold Pentylentetrazole induced seizure, Models for anxiolytic study i.e. Elevated plus maze, Models for nootropic study i.e. Object recognition test, Models for antidepressant study i.e. Forced swim test, Tail suspension test, Models for Muscle relaxant study i.e. Grip-strength test, The ethanol extract significantly and in dose dependent manner reduce the nociception induced by acetic acid. From the present study it was concluded that the herbal drugs can be potentially used to control the state of CNS disorders. Further investigations are, however, necessary to explore mechanism(s) of action involved in these pharmacological activities.

Keywords: *Melaleuca Leucadendra*, Neurobehavioural parameters, Epilepsy, Anxiolytic, Muscle relaxant study.

1. INTRODUCTION

Indian traditional medicine is predicated on phytochemical, pharmacological & allied approaches including instrumental techniques like chromatography, microscopy and others. There is accumulating evidence suggesting medicinal plants are illimitable reservoirs of drugs. Researchers with interest in natural products have intensified their effort to towards scientific evaluation of traditional medicines. There is accumulating evidence suggesting medicinal plants are illimitable reservoirs of drugs. The astonishing structural diversity among their active components makes them a subsidiary source of novel therapeutic compounds India has an antediluvian heritage of traditional medicine. (Mukherjee. et al., 1999) India has an antediluvian heritage of traditional medicine. Materica medica of India provides lots of information on the folklore practices and traditional aspects of therapeutically consequential natural product .Indian traditional medicine is predicated on phytochemical, pharmacological & allied approaches including instrumental techniques like chromatography, microscopy and others. There is accumulating evidence suggesting medicinal plants are illimitable reservoirs of drugs.

Recently available antipsychotics drugs are associated with variety of endocrine, allergic, hematopoietic, autonomic and neurological side effects. So, there is high prevalence of usage of traditional and alternative system of medicines for treatment of psychiatric disorders. In the search for new valuable products for the treatment of neurological disorders, medicinal plant research, worldwide, has progressed constantly, demonstrating the pharmacological effectiveness of different plant species in a variety of animal models. (Zhang ZJ. Et al.,2004) *Melaleuca Leucadendra* is traditionally used to reduce CNS disorders. Generally, plants possess many pharmacological actions, such as Hepatoprotective, antioxidant, anti-inflammatory, anthelmintic, antimicrobial since they contain numerous constituents of active chemicals in it Based on the claim by traditional healers that the plant is effective in the treatment of central nervous system (CNS) diseases. The present project is done to explore the potential of herbal drugs for the treatment of CNS disorders with a view to perform phytochemical investigation and assess Neurobehavioural and Neurochemical screening.

2. EXPERIMENTAL

2.1 Plant material

Plants materials *Melaleuca Leucadendra* bark were collected from the local market of Bhopal, (M.P.) during the month of May –July, 2022. The specimens were identified and authenticated by Dr. Zia ul Hassan, Assistant professor, Department of Botany, Saifia College of Science & Education, Bhopal and their herbarium was deposited. These collected specimens were chosen for the extraction process and assessment of Neurobehavioural activity.

2.2 Extraction

2.2.1 Ethanolic Extraction

The plant materials so collected were cleaned properly and washed with distilled water to remove dust particles and dried in shade. The dried drugs were coarsely powdered and then exhaustively extracted with 50% ethanol in Soxhlet apparatus for 72 h. The ethanolic extracts so obtained were freed of solvent under vacuum. (Yield: 9.33 %)

2.2.3 Qualitative analysis

The Ethanol extracts were screened for the presence of secondary metabolites. Tests were carried out for carbohydrates, reducing sugars, tannins, polyphenols, lipids, flavonoids, ketones, alkaloids, steroids and triterpenes.

3. IN VIVO STUDY

3.1 Animals for experiment

Acute toxicity studies were conducted by using albino mice of either sex weighing between 20 and 25 gms and healthy adult male albino rats weighing between 150 and 200 gms were selected for the antiurolithiatic activity. The animals were acclimatized to standard laboratory conditions (temperature: 25±20°C) and maintained on 12-h light: 12-dark cycle. They were provided with regular rat chow (Lipton India Ltd., Mumbai, India) and drinking water *ad libitum*.

3.2 Acute toxicity

In an acute toxicity study, animals were given single doses of drug. The Swiss albino rats were divided into groups. All animals fed with standard rat pelleted diet (Lipton India Ltd. pellets) and had free access to tap water *ad libitum*. Acute toxicity studies were performed according to the OECD guidelines. The doses selected for the study were 50 mg/kg, 100 mg/kg, 200 mg/kg, 300 mg/kg, 400 mg/kg for one day. Three animals were taken for each dose. It was observed that the extract didn't produce any significant effect on the behaviour of rats. The animals were observed for 3 hours after dose administration and also after 24 and 48 hours.

3.3 Screening for Neurobehavioural and Neurochemical screening

Screening for Neurobehavioural and Neurochemical screening was carried out in Wistar albino mice of either sex weighing 100-120 g.

3.3.1 Assessment of Epileptic activity

Wistar albino mice (25-35 g) bred in Central Animal facility of the Institute, were used. They were housed under standard conditions, maintained on a 12 h light/dark cycle and had free access to food and water up to the time of experimentation. The mice were acclimatized to the laboratory environment 1 h before the experiments. All experiments were conducted during the light period (08.00-16.00 h). During the experiments animals were free access to water only. All the protocols were approved by the Institutional Animal Ethical Committee (IAEC) and conducted according to the guidelines of CPCSEA (Committee for the Purpose of Control and Supervision of Experiments on Animals).

3.3.2 Pentylenetetrazol (PTZ)-induced convulsion test

Animal groups

Four groups of mice (n = 10) were used. Group I was administered the vehicle, i.e., normal saline (1 ml/100g body weight) and served as control, Group II received reference standard (diazepam, 2 mg/kg, i.p.) while Groups III and IV were administered extract 100 and 200 mg/kg, p.o., respectively, of the extracts. Two hours later, PTZ was administered (60 mg/kg, i.p.) to all four groups. The animals were observed for 30 min and the onset and duration of convulsion noted. (Kitano. et al., 2005)

3.3.3 Assessment of anxiolytic study

Animal groups

Elevated plus – Maze test:

Animals were randomly allocated to four experimental groups (n= 5 each). Group 1 and 2 were named as negative and positive control and 2 other groups were termed as treated group. (Dhananjaya. et al., 2011) The test groups received ethanolic extract of *Melaleuca Leucadendra* at the dose of 200 and 400mg/kg body weight respectively. Group 1 received 1% Tween 80 solution. Group 2 got administration of Diazepam (as a standard drug) at 1mg/kg bodyweight. Drug or vehicle was injected intraperitoneally in a volume of 0.5ml/kg. Tests were performed 30 min after injections. (Navarro JF et al., 2006).

3.3.4 Assessment of nootropic study

Object recognition test

A plastic chamber (35cm×35cm×35 cm) was used in low light condition (about 40 lx) during the light phase of the light/dark cycle. The general procedure, as described elsewhere, consisted of three different phases: a habituation phase, an acquisition phase, and a retention phase. On the 1st day (habituation phase), mice were individually subjected to a single familiarization session of 10 min, during which they were introduced in the empty arena, in order to become familiar with the apparatus. On the 2nd day (acquisition phase) animals were subjected to a single 10-min

session, during which floor-fixed two objects (A and B) were placed in a symmetric position in the central line of the arena, 10cm from each and 8 cm from the nearest wall (each object occupies approximately 5 cm space by its size). The two objects, made of the same wooden material with the similar color and smell, were different in shape but identical in size. Mice were allowed to explore the objects in the open field. The exploration time on each object was shown (as seconds) to indicate the exploring activity of mice. On the 3rd day (retention phase), mice were allowed to explore the open field in the presence of two objects: the familiar object A and a novel object C in different shapes but in similar color and size (A and C). A recognition index (for retention session), calculated for each mouse, was expressed as the ratio $(TC \times 100) / (TA + TC)$, where TA and TC are the time spent during retention phase on object A and object C, respectively. The time spent exploring any object (nose pointing toward the object at a distance ≤ 1 cm, but not mounting on the object or playing with the object) was recorded (using stopwatch) by hand. (Shete. et al., 2010)

3.3.5 Assessment of antidepressant study

Forced Swimming Test (FST)

Mice of either sex were individually forced to swim in an open cylindrical container (diameter 10 cm, height 25 cm), containing 19 cm of water at $25 \pm 1^\circ\text{C}$. The immobility time, defined as the absence of escape-oriented behaviors, such as swimming, was scored during 6min with the help of stop-watch, as described previously by (Eckeli *et al.*, 2000). All the mice of either sex were divided in five different groups. The first group assigned as control receiving only vehicle (NaCl 5 mL kg⁻¹). The other groups received acute dose of extracts. The total duration of immobility was recorded during the last 6 min of the 10 min period, where the activity in the two first minutes is discarded (Porsolt, 1981). Each mouse was judged to be immobile when it ceased struggling and remained floating motionless in the water, making only those movements necessary to keep its head above water. A decrease in the duration of immobility is indicative of an antidepressant like effect. For the next exposure of crude extracts, FST of repetitive doses of crude extracts were assessed after 3 days of treatment within 30 min after the last dose of administration. During the test session, the immobility time was recorded. The mice were considered immobile when neither hind leg was moving; the mice were slightly hunched forward. Another reason for choosing this animal model is the correlation which is observed between results in this model and clinical potency, which is not found in any other models (Willner, 1984; Shah et al., 2006). A different interpretation of the FST holds that rats or mice, in this behavioral paradigm, learn to be immobile. Immobility is considered to be an adaptive response to the situation consisting of animals learning to keep their heads out of water with a minimum of energy expenditure (Parra *et al.*, 1999).

Tail Suspension Test (TST)

The total duration of immobility induced by tail suspension test was measured according to the method described by Steru *et al.* (1985). Mice both acoustically and visually isolated were suspended 70 cm above the floor by adhesive tape placed approximately 1 cm from the tip of the tail. The total immobility period was scored manually during 6 min test session with the help of stopwatch. Immobility was defined as the absence of any limb or body movements, except for those caused by respiration or when they hung passively and completely motionless. The parameter obtained was the number of seconds spent immobile. Parameter used was the number of seconds spent immobile. (Cryan. et al., 2005)

3.3.6 Assessment of Muscle relaxant study

Grip strength test:

A Rota-rod (Inco- Ambala, Instruments and chemicals Pvt. Ltd. Model town, Ambala City -03) was used to measure the grip strength in mice. The instrument (a horizontal rotation device) was set at a rate of 16 revolutions per minute^{5,6}. The animals were placed on the rotating rod and fall off time i.e, when the animal falls from the rotating rod, was recorded, which was taken as grip strength. Diazepam (26 mg/kg i.p.) was used as the standard drug. Grip strength in all the groups was measured before and at 30 minutes, 1 h, 2 h, 4 h, and 6 h of the administration of the dose. (Rogers. et al., 1997)

4. RESULTS AND DISCUSSION

4.1 Qualitative analysis

The preliminary phytochemical screening of extracts revealed the presence of various phyto-constituents. The ethanolic extract of *Melaleuca Leucadendra* revealed the presence of alkaloids, glycosides, sterols, flavonoids, tannins, phenolic compounds, carbohydrates, fats and oils.

4.2.1 Pentylenetetrazol (PTZ)-induced convulsion test

On the basis of phytochemical investigations, the pharmacological screening was performed on ethanolic extracts of selected plants on Pentylenetetrazol (PTZ)-induced convulsion test. Table 6.5 shows that the extract increased the onset of pentylenetetrazol-induced seizures but reduced the duration of the convulsions. The increase in the onset of seizure was statistically significant, $p < 0.01$ and $p < 0.05$ for extract doses of 100 and 200 mg/kg, respectively, and highly significant ($p < 0.001$) for the standard drug, diazepam; with regard to their duration reduction effect, the difference was statistically significant at $p < 0.01$ for both doses. However, the clonus phase was completely abolished only at the lower dose of the extract and also with diazepam.

Table 1: Effect of *Melaleuca Leucadendra* on pentylenetetrazol (PTZ)-induced seizures in mice

Groups	Onset of convulsion (min)	Duration of convulsion (min)
Vehicle(control)	1.01 \pm 0.11	38.2 \pm 0.2
Standard (Diazepam)	0 \pm 0 [†]	0 \pm 0 [†]
Etoh (100mg/kg)	5.94 \pm 0.17 [#]	6.9 \pm 0.1 [#]
Etoh (200mg/kg)	1.59 \pm 0.13 [*]	16.1 \pm 0.1 [#]

[†]P < 0.001; [#]p < 0.01; ^{*}p < 0.05




Figure 1: Effect of *Melaleuca Leucadendra* on pentylenetetrazol (PTZ)-induced seizures

4.2.2 Assessment of anxiolytic study

Elevated plus – Maze test:

From the experiment it was observed that mice taken ethanol at dose of 200 and 400 mg/kg body weight, stayed more time in open arm of Elevated plus Maze apparatus in comparison to standard and negative control group. Moreover they were also stayed less time in closed arm of Elevated plus Maze apparatus in comparison to standard and negative control group. The value obtained from these extract were statistically significant ($p < 0.05$). From the observed result, it could be concluded that ethanolic extract may have anxiolytic activity.

Table 2: Effect of *Melaleuca Leucadendra* on Elevated plus Maze experiment

Mean \pm SD (counts/5minutes)

Animal Group	Open arm Duration	Frequency of open arm	Closed arm Duration	Frequency of Closed arm
Vehicle(control)	2.6	0.8	257.8 \pm 11.39398	8.4 \pm 1.719
Standard (Diazepam)	2.4	0.2	261.8 \pm 11.39398	7.6 \pm 1.719
Etoh (200mg/kg)	16	3.2	207.6 \pm 11.39398	13.2 \pm 1.719
Etoh (400mg/kg)	8.2	2.6	203.2 \pm 11.39398	14.4 \pm 1.719




Figure 2: Average time duration in different arms and their frequency in Elevated Plus

4.2.3 Assessment of nootropic study

Object recognition test

The increase in Recognition Index (RI) indicates nootropic effect. The RI of control group was 51.58 ± 1.8 %. Scopolamine (0.3mg/kg) shown significant ($p < 0.05$) decrease in the RI to 30.60 ± 3.2 %. Piracetam (100mg/kg) has significantly ($p < 0.01$) increased the RI to 73.8913 ± 2.6 % and also exhibited significant ($p < 0.01$) antagonism of the amnesic effect of scopolamine. The extract in the doses 10 mg/kg increased the RI to 53.70 ± 3.6 % ($p < 0.05$) respectively. Table 6.7

Table 3: Effect of *Melaleuca Leucadendra* on object recognition test

Animal Group (Dose in mg/kg)	Trial-1	Trial-2 (sec)		Recognition Index
	Time spent (sec)	Time spent familiar object	Time spent new object	
Control	20.00 \pm 3.05	10.10 \pm 0.47	11.10 \pm 1.27	51.58 \pm 1.84
Scopolamine(0.3)	31.11 \pm 3.84	31.98 \pm 20.98**	14.45 \pm 2.48	30.60 \pm 3.24*
Piracetam (100)	36.80 \pm 4.04	15.80 \pm 2.60	42.80 \pm 2.27**	73.89 \pm 2.65**
Ethanolic (10)	22.46 \pm 5.09	20.45 \pm 2.96	24.30 \pm 4.58	53.70 \pm 3.69




Figure 3: Effect of *Melaleuca Leucadendra* on object recognition test

4.2.4 Assessment of antidepressant study

Forced Swimming Test (FST)

According to FST results from the first day was shown in Table 6.8. A significant ($p < 0.05$) decrease in the duration of immobility was seen with two standard drug haloperidol and fluoxetine and The results obtained after a single administration of the extract suspension showed that the immobility time of mice decreased dose dependently which mice were more active in both employed models and it means that the antidepressant effect was stronger. However, for all two doses administered there were differences compared to the control, that is, they led to reduction of immobility time.

Table 4: Effect of *Melaleuca Leucadendra* Forced Swimming Test (FST)

Animal Group (Dose in mg/kg)	Immobility time (s) at first day	Change (%)	Immobility time (s) at third day	Change (%)
Control(0)	386.0 ± 31.5	-	385.0 ± 52.8	-
Haloperidol(1)	115.7 ± 4.5***	-71.84	74.8 ± 8.2***	-80.62
Fluoxetine(10)	211.7 ± 6.1***	-46.25	200.0 ± 7.0***	-48.06
ET(50)	380.3 ± 21.7ns	0.79	235.3 ± 41.8*	-38.17




Figure 4: Effect of *Melaleuca Leucadendra* Forced Swimming Test (FST)

Table 5: Effect of *Melaleuca Leucadendra* Tail suspension test (TST)

Animal Group (Dose in mg/kg)	Immobility time (s) at first day	Change (%)
Control(0)	315.3 ± 30.4	42.39
Haloperidol(1)	61.8 ± 15.1***	-80.39
Fluoxetine(10)	91.8 ± 11.5***	-69.97
ET(50)	177.0 ± 9.5***	-42.39

a Mean±SD, n = 3; Statistically significant difference compared to the control group of animals: *** p<0.05, ** p<0.01, * p<0.001 and ns = Data are not different significant compared to control (vehicle)




Figure 5: Effect of *Melaleuca Leucadendra* Tail suspension test (TST)

4.2.5 Assessment of Muscle relaxant study

Grip Strength Test

Grip strength in seconds

Group	Treatment	Dose (mg/kg)	0 hr	30 min	1 hr	2 hr	4 hr	6 hr
1	Control	--	344.17 ± 11.620	359.33 ± 19.416	328.23 ± 12.456	340.50 ± 12.132	376.33 ± 11.071	360.17 ± 19.511
2	Diazepam	26	359.50 ± 18.936	* 221.17 ± 13.893	* 124.17 ± 13.946	** 124.23 ± 11.923	** 100 ± 4.310	** 102 ± 4.310
3	Ethanollic extract	200	341.50 ± 12.233	* 318.17 ± 10.778	* 335.23 ± 9.127	ns 360.01 ± 11.923	358.50 ± 10.987	337.23 ± 6.940

Values are mean ± SME; n=6 in each group; *significantly different at $p < 0.05$; ** significantly different at $p < 0.01$; ns- Non.significant

Table 6: Effect of *Melaleuca Leucadendra* on Grip Strength Test




Figure 6: Effect of *Melaleuca Leucadendra* on Grip Strength Test

5. CONCLUSION

The preliminary phytochemical screening of extracts revealed the presence of various phyto-constituents. The ethanollic extract of *Melaleuca Leucadendra* revealed the presence of alkaloids, glycosides, sterols, flavonoids, tannins, phenolic compounds, carbohydrates, fats and oils. The pharmacological screening was performed on ethanollic extracts of selected plants to confirm the Neurobehavioural and Neurochemical screening of *Melaleuca Leucadendra* ethanollic extract the extract increased the onset of pentylenetetrazol-induced seizures but reduced the duration of the convulsions. The increase in the onset of seizure was statistically significant, $p < 0.01$ and $p < 0.05$ for extract doses of 100 and 200 mg/kg, respectively, and highly significant ($p < 0.001$) for the standard drug, diazepam; with regard to their duration reduction effect, the difference was statistically significant at $p < 0.01$ for both doses. However, the clonus phase was completely abolished only at the lower dose of the extract and also with diazepam.

From the experiment it was observed that mice taken ethanol at dose of 200 and 400 mg/kg body weight, stayed more time in open arm of Elevated Plus Maze apparatus in comparison to standard and negative control group. Moreover they were also stayed less time in closed arm of Elevated plus Maze apparatus in comparison to standard and negative

control group. The value obtained from these extract were statistically significant ($p < 0.05$). From the observed result, it could be concluded that ethanolic extract may have anxiolytic activity.

The increase in Recognition Index (RI) indicates nootropic effect. The RI of control group was 51.58 ± 1.8 %. Scopolamine (0.3mg/kg) shown significant ($p < 0.05$) decrease in the RI to 30.60 ± 3.2 %. Piracetam (100mg/kg) has significantly ($p < 0.01$) increased the RI to 73.89 ± 2.6 % and also exhibited significant ($p < 0.01$) antagonism of the amnesic effect of scopolamine. The BF in the doses 10 mg/kg increased the RI to 53.70 ± 3.6 % ($p < 0.05$) respectively.

According to FST results from the first day was shown in Table 6.8 A significant ($p < 0.05$) decrease in the duration of immobility was seen with two standard drug haloperidol and fluoxetine and The results obtained after a single administration of the extract suspension showed that the immobility time of mice decreased dose dependently which mice were more active in both employed models and it means that the antidepressant effect was stronger. However, for all two doses administered there were differences compared to the control, that is, they led to reduction of immobility time. However, ET exhibited highest decrease of duration immobility time in TST (77.61%) followed by haloperidol as a positive control (80.62%).

6. ACKNOWLEDGEMENT

I am thankful to my mentor and my research institute for research Project.

7. REFERENCES

- [1] Abramson, Modeling psychopathology in the laboratory, *Psychopathology: Experimental Models*. San Francisco (1978), 1-26.
- [2] Adamec, Long-lasting, selective, anxiogenic effects. *Physiology and Behaviour* (2004), 83, 401-410.
- [3] Akiskal, Interaction of biologic and psychologic factors. *Acta Psychiatrica Scandium* (1985), 71, 131-139.
- [4] Alberts, Olfactory bulb removal. *Nature* (1972), 238, 25-28.
- [5] Alonso, Population level. *British Journal Psychiatry* (2007), 190, 299-306.
- [6] American Psychiatric Association. In: Text revision. Fourth ed. Washington, DC: *American Psychiatric Association* (2000), 780-783.
- [7] American Psychiatric Association. DSM-V, Co morbidity of Depression and Generalized Anxiety Disorder (2007) Washington, DC: *American Psychiatric Association*, 320-324.
- [8] Andrade, Longitudinal study of daily variation of rats, *Physiology and Behaviour* (2003), 78, 125-133.
- [9] Anderson, Treatment discontinuation with selective serotonin reuptake inhibitors. *British Medical Journal* (1995), 310, 1433-1438.
- [10] Andersen, Behavioural evaluation of methods. *Applied Animal Behaviour Science* (2000), 69, 227-240.
- [11] Andreatini, The relationship between anxiety and depression. *Brazilian Journal of Medical and Biological Research* (1999), 32, 1121-1126.
- [12] Abdel-Waheb. Investigation of the flavanoid content of *Salvadora persica* L. *Bull Fac Pharm* (1990), 28, 67-70.
- [13] Adebisi A, Modulation of jejunal contractions. *Phyto Res* (2005), 19, 628-63.
- [14] Adeneye AA. Preliminary hypoglycemic and hypolipidemic activities. *Biol Med* (2009), 1, 1-10.
- [15] Adeyemi DO. Anti hyperlipidemic activities of *Annona muricata*. *Int J Alter Med* (2009), 7, 62-67.
- [16] Agarwal N, Behavioural and lethal effects of alcoholic extracts of *Evolvulus alsinoides* in albino mice. *J Physio Allied Sci* (1997), 31, 81.
- [17] Ahmad S. Anticonvulsant potential of callus cultures. *Ori Pharm Exp Med* (2007), 7, 46-50.
- [18] Alali F. GC-MS analysis and bioactivity testing of the volatile oil. *Nat Prod Res* (2003), 17, 189-194.
- [19] Al-Bagieh. Antiherpes simplex c-virus type 1 activity. *Biomed Lett* (1990), 47, 67-70.
- [20] Alder R. A systematic review of the effectiveness of garlic. *J Am Acad Nurse Pract* (2003), 15, 120-9.
- [21] Al-Howiriny T. Gastric antiulcer of celery (*Apium graveolens*) in rats. *Pharm Biol*, 48, 786-793.
- [22] Berlyne. Novelty, arousal, and the reinforcement. *Journal of Comparative and Physiology Psychology* (1966), 62, 222-226.
- [23] Berman. Antidepressant effects of ketamine in depressed patients. *Biological Psychiatry* (2000), 47,

351-354.

- [24] Berrios. Conceptual issues. P, editors. *Biological Psychiatry* (2002). John Willey & Sons, 1-24.
- [25] Bhattacharya. Antioxidant effect of Withania somnifera glycowithanolides. *Journal of Ethanopharmacology* (2001), 74, 1-6.
- [26] Bill. Current and residual functional disability: The Netherlands Mental Health Survey and Incidence Study (NEM-ESIS). *Psychological Medicine* (2000), 30, 657-668.
- [27] Blanchard. The mouse defense test battery: pharmacological and behavioral assays for anxiety and panic. *European Journal of Pharmacology* (2003), 463, 97-116.
- [28] Blazer, Social support and mortality in an elderly community population. *American Journal of Epidemiology* (1982), 115, 684-694.
- [29] Blier, Current advances and trends in the treatment of depression. *Trends in Pharmacological Science* (1994), 15, 220-226.
- [30] Borsini, Is the forced swimming test a suitable model for revealing anti-depressant activity? *Psychopharmacology* (1988), 94, 147-160.
- [31] Choubey Ankur. Evaluation of the immunomodulatory activity of methanolic and ethanolic extract of leaves of aegle marmelos in rats, *International Journal of Drug Development & Research*, July-Sept (2010), Vol. 2(3)
- [32] Clement Y. Biological Basis of Anxiety. *Neuroscience and Biobehavioural Reviews* (1998, 22(5): 623-633.
- [33] Cryan. Tail suspension test as a model for assessing antidepressant activity. *Neuroscience and Biobehavioral Reviews* (2005), 29 (571-625).
- [34] Dhananjaya D.R, (2011), Anxiolytic activity of ethanolic extract of Trigonella foenum graecum seeds. *Archives of Applied Science Research*, 3 (1): 91-95.
- [35] Baeshen HA. Uptake and release of fluoride from Fluoride-impregnated chewing sticks *in vitro* and *in vivo*. *Caries Res* (2008), 42, 363-8.
- [36] Bisset, Herbal Drugs and Phytopharmaceuticals. Stuttgart: Medpharm Scientific Publishers (1994), p. 70-72.
- [37] Carroll, A specific laboratory test for the diagnosis of melancholia: Standardization, validation and clinical utility. *Archives of General Psychiatry* (1981), 38, 15-22.
- [38] Carver, Relations of serotonin function to personality: Current views and a key methodological issue. *Psychiatry Research* (2006), 144, 1- 15.
- [39] Carver, Origins and functions of positive and negative affect: A control-process view. *Psychological Review* (1990), 97, 19-35.
- [40] Celada, The therapeutic role of 5-HT_{1A} and 5-HT_{2A} receptors in depression. *Journal of Psychiatry Neuroscience* (2004), 29, 252-265.
- [41] Charney. Depression and Bipolar Support. *Archives of General Psychiatry* (2003), 60, 664 -672.
- [42] Charney. Monoamine dysfunction treatment of depression. *Journal of Clinical Psychiatry* (1997), 59, 11-14.
- [43] Chrousos. The concepts of stress. *Journal of American Medical Association* (1992), 267, 1244-1252.
- [44] Clark. Psychometric evidence and psychometric implications. *Journal of Abnormal Psychology* (1991), 100, 316-336.
- [45] Coulehan. Medical comorbidity. *Archives of Internal Medicine* (1990), 150, 2363-2367.
- [46] Crawley. Preliminary report of a simple animal behavior. *Pharmacology Biochemistry and Behavior* (1980), 13, 167-170.
- [47] Crawley. Behavioural phenotyping of transgenic and knockout mice. NY: Wiley-Liss (2000), 386-448.
- [48] Crowley, J.J., Lucki, I. (2005) Opportunities to discover genes regulating depression and antidepressant response from rodent behavioural genetics. *Current Pharmaceutical Design*, 11, 157-169.
- [49] Cryan. Onset of the effects of the 5-HT_{1A} antagonist, WAY-100635. *Pharmacology Biochemistry and Behavior* (1999), 63, 333-338.
- [50] Cryan. Assessing substrates underlying the behavioral effects of antidepressants. *Neuroscience*

- Biobehavioral Review* (2005), 29, 547–569.
- [51] Cryan. In search of a depressed mouse. *Molecular Psychiatry* (2004), 9, 326–357.
- [52] Cryan. Assessing antidepressant activity in rodents: recent developments and future needs. *Trends in Pharmacological Science* (2002), 23, 238-245.
- [53] Dam. The sero-otonin transporter and 5HT_{2A} receptor lesions. *Neurological Research* (2007), 29, 717-722.
- [54] Dantzer. Cytokine, sickness behavior, and depression. *Clinical Neurology* (2006), 24, 441–460.
- [55] Davidson. Achieving remission with venlafaxine and fluoxetine. *Depression Anxiety* (2002), 16, 4–13.
- [56] Davis. Long-lasting effects of acute stress on the neurochemistry. *Psychopharmacology*, 118, 267– 272.
- [57] Deakin. Roles of serotonergic systems in escape, *Psychopharmacology Vol 2*. New York: Academic Press (1983), 149-193.
- [58] Deakin. Distinct roles of 5HT subsystems in panic, anxiety and depression. (Eds.), *Biol. Psych.*, vol. 1. Elsevier, Amsterdam (1991), 305–307.
- [59] Deakin. 5-HT, anti-depressant drugs and the psychosocial origins of depression. *Journal of Psychopharmacology* (1996), 10, 31–38.
- [60] Deakin. Social stress, steroid levels, 5-HT and depression. *Journal of Psychopharmacology* (1997), 11, A93-A97.
- [61] Deakin. Depression and antisocial personality disorder. *Journal of Neural Transmission Supplementum* (2003), 64, 79–93.
- [62] Deb.. Rate of psychiatric illness one year after traumatic brain injury. *American Journal of Psychiatry* (1999), 156, 374 –378.
- [63] Denenberg, V.H. (1969) Open-field behavior in the rat: what does mean? *Annals of the Newyork Academy of Science*, 159, 852–859.
- [64] Edwards, The psychological impact of a cancer diagnosis on families. *Psycho-oncology* (2004), 13, 562-576.
- [65] Edwards. Olfactory bulb removal: effects on sexual behaviour and partner-preference in male rats. *Physiology and Behaviour* (1990), 48, 447–450.
- [66] Evans. Mood disorders in the medically ill. *Biological Psychiatry* (2005), 58, 175-189.
- [67] Fann, Sertraline in the treatment of major depression. *Journal of Neuro-psychiatry Clinical Neuroscience* (2000), 12, 226–32.
- [68] Fedoroff, Depression in patients with acute traumatic brain injury. *American Journal of Psychiatry* (1992), 149, 918-923.
- [69] Feinstein, The pre-therapeutic classification of comorbidity in chronic disease. *Journal of Chronic Diseases* (1970), 23, 455-468.
- [70] Ferré. Struggling and flumazenil effects in the swimming test are related to the level of anxiety in mice. *Neuropsychobiology* (1994), 29, 23-27.
- [71] Fehr. Serotonergic polymorphisms in patients suffering from alcoholism. *Progress in Neuropsychopharmacology and Biology Psychiatr* (2001) y, 25, 965–982.
- [72] Fernandez-Ruiz. Olfactory dysfunction in hereditary ataxia and basal ganglia disorders. *Neuroreports* (2003), 14, 1339–1341.
- [73] Feske. Anxiety as a predictor of response to interpersonal psychotherapy. *Depression and Anxiety* (1998), 8, 135–141.
- [74] Field. Pregnancy anxiety and comorbid depression. *Depression and Anxiety* (2003) 17, 140–151.
- [75] File. Can social interaction be used to measure anxiety *British Journal of Pharmacology* (1978), 62, 19–24.